Quantitative Methods: Reliability, Validity, and Use of SEM to Assess Psychometric Equivalence

> Ron D. Hays, Ph.D. UCLA Department of Medicine Los Angeles, CA

> > November 18, 2005

Measurement Range for Health Outcome Measures



Indicators of Acceptability

Response rate
Administration time
Missing data (item, scale)

Variability

 All scale levels are represented
 Distribution approximates bell-shaped "normal"



Measurement Error

Observed = true + systematic + random score error error



Flavors of Reliability

Test-retest (administrators)
Intra-rater (raters)
Internal consistency (items)

Intraclass Correlation and Reliability

Model	<u>Reliability</u>	Intraclass Correlation
One-Way	$\frac{MS_{BMS} - MS_{WMS}}{MS_{BMS}}$	$\frac{MS_{BMS} - MS_{WMS}}{MS_{BMS} + (K-1)MS_{WMS}}$
Two-Way Fixed	$\frac{MS_{BMS} - MS_{EMS}}{MS_{BMS}}$	$\frac{MS_{BMS} - MS_{EMS}}{MS_{EMS} + (K - 1)MS_{EMS}}$
Two-Way	$N(MS_{BMS} - MS_{EMS})$	$MS_{\rm DMG} - MS_{\rm DMG}$

Random

 $\frac{N(MS_{BMS} - MS_{EMS})}{NMS_{BMS} + MS_{JMS} - MS_{EMS}}$

 $\frac{MS_{BMS} - MS_{EMS}}{MS_{BMS} + (K-1)MS_{EMS} + K(MS_{JMS} - MS_{EMS})/N}$

Cronbach's Alpha

Source	<u>df</u>	<u>SS</u>	<u>MS</u>
Respondents (BMS)	4	11.6	2.9
Items (JMS)	1	0.1	0.1
Resp. x Items (EMS)	4	4.4	1.1
Total	9	16.1	

Alpha =
$$\frac{2.9 - 1.1}{2.9} = \frac{1.8}{2.9} = 0.62$$

Reliability Minimum Standards

0.70 or above (for group comparisons)
0.90 or higher (for individual assessment)

 \gg SEM = SD (1- reliability)^{1/2}

Construct Validity

Does measure relate to other measures in ways consistent with hypothesis?

Responsiveness to change

Responsiveness to Change and Minimally Important Difference

 HRQOL measures should be responsive to interventions that changes HRQOL

 Evaluating responsiveness requires assessment of HRQOL

» pre-post intervention of known efficacy

- » at two times in tandem with anchor
- » HRQOL change among people who changed on anchor

Self-Report Anchor

• Overall has there been any change in your asthma since the beginning of the study? Much improved; Moderately improved; Minimally improved No change Much worse; Moderately worse; Minimally worse

Clinical Anchor

"changed" group = seizure free (100% reduction in seizure frequency)
"unchanged" group = < 50% change in seizure frequency

Responsiveness Indices

- (1) Effect size (ES) = D/SD
- (2) Standardized Response Mean $(SRM) = D/SD^{\dagger}$
- (3) Guyatt responsiveness statistic (RS) = D/SD^{\ddagger}
- D = raw score change in "changed" group; SD = baseline SD;
 - $^{\dagger}SD = SD \text{ of } D;$
 - \$ SD = SD of D among "unchanged"

Effect Size Benchmarks

Small: 0.20->0.49
Moderate: 0.50->0.79
Large: 0.80 or above



Hypothetical Multitrait/Multi-Item Correlation Matrix

	<u>Trait #1</u>	<u>Trait #2</u>	<u>Trait #3</u>
Item #1	0.80*	0.20	0.20
Item #2	0.80*	0.20	0.20
Item #3	0.80*	0.20	0.20
Item #4	0.20	0.80*	0.20
Item #5	0.20	0.80*	0.20
Item #6	0.20	0.80*	0.20
Item #7	0.20	0.20	0.80*
Item #8	0.20	0.20	0.80*
Item #9	0.20	0.20	0.80*

*Item-scale correlation, corrected for overlap.

Confirmatory Factor Analysis

- Compares observed covariances with covariances generated by hypothesized model
- Statistical and practical tests of fit
- Factor loadings
- Correlations between factors
- Regression coefficients

Fit Indices



Parameter Estimates (standardized) for Confirmatory Factor Analytic Model Estimated Separately Among English and Spanish (in parentheses) Respondents

	Physical Health	Mental Health	Evaluations of Care
Physical functioning	0.69 (0.25)		
Role-physical	0.81 (0.53)		
Pain	0.69 (0.88)	$0.12(0.02^3)$	
General health perceptions	0.56 (0.39)	0.28 (0.34)	
Emotional well-being		0.86 (0.93)	
Role-emotional		0.62 (0.44)	
Energy/fatigue	0.38 (0.22)	0.56 (0.69)	
Social functioning	0.41 (0.45)	0.35 (0.36)	
Doctor			0.96 (0.97)
Overall			0.93 (0.92)
Access			0.81 (0.80)
Wait time			0.75 (0.79)
Choice			0.73 (0.78)
Plan			0.70 (0.70)

NOTE: Estimated correlations for English and Spanish respondents, respectively, were as follows: physical and mental health (r = .45, .55), mental health and patient evaluations of care (r = .23, .31), and physical health and patient evaluations of care ($r = .05, .16^{a}$). a. Not statistically significant.

Parameter Estimates (standardized) for Confirmatory Factor Analytic Model Constraining Factor Loadings and Correlations to Be Equal for English and Spanish Respondents

	Physical Health	Mental Health	Evaluations of Care
Physical functioning	0.69 (0.51)		
Role-physical	0.81 (0.78)		
Pain	0.69 (0.68)	0.12 (0.12)	
General health perceptions	0.56 (0.53)	0.28 (0.27)	
Emotional well-being		0.86 (0.80)	
Role-emotional		0.62 (0.57)	
Energy/fatigue	0.37 (0.37)	0.56 (0.56)	
Social functioning	0.41 (0.48)	0.35 (0.41)	
Doctor			0.96 (0.97)
Overall			0.93 (0.93)
Access			0.81 (0.81)
Wait time			0.75 (0.80)
Choice			0.73 (0.75)
Plan			0.70 (0.74)

NOTE: Estimated correlations for English and Spanish respondents, respectively, were as follows: physical and mental health (r = .45), mental health and patient evaluations of care (r = .23), and physical health and patient evaluations of care (r = .06).

Acknowledgments

 Supported in whole by UCLA Center for Health Improvement in Minority Elders/ Resource Centers for Minority Aging Research, National Institute on Aging (AG-02-004)