Appendix C
T Scores Obtained by Cumulative Percentages of the Normative Samples

T Scores Obtained by Cumulative Percentages of the Normative Samples: Physical Functioning Scale (PF) Table C.I.

| T | | Age-Base | d Sample | | Age-S | Stratified S | ample | τ |
|-------|------|----------|----------|-----------------|--------------|--------------|-------|-------|
| Score | 1824 | 25-44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | l |
| .2 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 2 |
| | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 3 |
| 4 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | -4 |
| 5 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 5 |
| - 6 | 11.5 | <0.1 | <0.1 | <0.1 | ≺ 0.1 | <0.1 | <0.1 | - 6 |
| 7 | 2.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 7 |
| . 8 | 2.0 | 20.5 | <0.1 | <0.1 | <0,1 | <0.1 | <0.1 | . 8 |
| 9 | 2.0 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 9 |
| 10 | 2.0 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 10 |
| II | 2.0 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | |
| 12 | 2.0 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 12 |
| 13 | 2.0 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 13 |
| 14 | 2.0 | 1:0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.i | 14 |
| 15 | 2.0 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 15 |
| 16 | 2.0 | 1.0 | <0.1 | < 0.1 | 0.6 | <0.1 | 0.4 | 16 |
| 17 | 2.0 | 2.0 | <0.1 | <0.i | 0.6 | 0.8 | 0.8 | 17 |
| . 18 | 2.0 | 2.5 | <0.1 | <0.1 | 0.8 | 0.8 | 1.2 | 18 |
| 19 | 2.0 | 2.5 | <0.1 | <0.1 | 1.0 | 0.8 | 1.2 | 19 |
| 20 | 2.0 | 2.5 | <0.1 | <0.1 | 1.0 | 8.0 | 1.2 | 20 |
| . 21 | 3.0 | 3.0 | <0.1 | <0.1 | 1.4 | 0.8 | 1.6 | 21 |
| .22 | 3:0 | 3,0 | 0.5 | <0.1 | 2.0 | 2.0 | 2.0 | 22 |
| 23 | 3.0 | 3.5 | 1.0 | <0.1 | 2.4 | 2.4 | 2.0 | 23 |
| -24 | 3.5 | 3.5 | 2:0 | 1.0 | .2.8 | 3:5 | 2.9 | 24 |
| 25 | 3.5 | 5.5 | 2.5 | 1.0 | 3.6 | 3.9 | 4.5 | 25 |
| 26 | 4.5 | 5.5 | 3.0 | 1.0 | 4.2 | 3.9 | 4:9 | 26 |

Table C.I. (Continued)

| Т | | Age-Based | l Sample | | Age-S | Stratified Sa | ample | T |
|-------|-------|-----------|----------|-------|---------|---------------|-------|-------|
| Score | 18-24 | 25-44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| 27 | 4.5 | 5.5 | 4.0 | 1.5 | 5.2 | 4.7 | 5.7 | 27 |
| 28 | 5.0 | 6.5 | 5.0 | 2.0 | 6.0 | 5.5 | 6.9 | 28 |
| 29 | 6.0 | 7.0 | 6.5 | 2.5 | 7.6 | 7.5 | 7.8 | 29 |
| 30 | 6.5 | 7.0 | 8.0 | 3.0 | 7.8 | 7.5 | 8.2 | 30 |
| 31 | 7.0 | 7.0 | 9.0 | 5.5 | 8.4 | 7.8 | 9.0 | 31 |
| 32 | 8.0 | 7.0 | 11:5 | 6.5 | 9,0 ≉ | 9.0 | 9.4 | 32 |
| 33 | 8.5 | 8.5 | 12.0 | 8.5 | 9.6 | 9.4 | 10.2 | 33 |
| 34 | 8.5 | 10.0 | 12.0 | 11.0 | 8.01 | 9.8 | 11.8 | 34 |
| 35 | 9.0 | 10.5 | 14.5 | 13.0 | 12.0 | 11.8 | 11.8 | 35 |
| 36 | 9.0 | 11.0 | 14.5 | 15.5 | 12.2 | 12.2 | 13.1 | 36 |
| 37 | 9.5 | 11.5 | 15.0 | 16.5 | 13.2 | 12.2 | 13.5 | 37 |
| 38. | 9.5 | 12.0 | 15.5 | 17.0 | 14.0 | 13.3 | 13.5 | 38 |
| 39 | 10.5 | 13.0 | 15.5 | 17.5 | 14.8 | 14.9 | 14.7 | 39 |
| 40 | 10.5 | 14.0 | 17.0 | 19.0 | 16.0 | 16.9 | 15.9 | 40 |
| 41 | 12.0 | 14.5 | 18.0 | 21.0 | 18.4 | 18.4 | 17.6 | 41 |
| -42 | 12.0 | 16.5 | 20.5 | .23.0 | 19.8 | 20.8 | 19.2 | -42 |
| 43 | 13.5 | 18.0 | 21.5 | 24.5 | 21.6 | 23.1 | 20.0 | 43 |
| -44 | 14.0 | 18.5 | 22.5 | 27.0 | 23.2 | 25.1 | 21.6 | 44 |
| 45 | 15.5 | 21.0 | 24.5 | 29.0 | 25.0 | 25.9 | 23.7 | 45 |
| 46 | 16.5 | 24.0 | 27.0 | 31.5 | 26.6 | 28.6 | 24.9 | 46 |
| 47 | 19.5 | 25.0 | 31.0 | 35.5 | 28.2 | 29.0 | 26.5 | 47 |
| 48 | 20.5 | 26:5 | 34.5 | 38.5 | 30.6 | 32.2 | 27.3 | 48 |
| 49 | 20.5 | 32.0 | 35.0 | 42.5 | 33.2 | 35.7 | 32.7 | 49 |
| .50 | .20.5 | 34.5 | 37.0 | 45.0 | 37.8 | 38.0 | 35.9 | 50 |
| 51 | 36.5 | 34.5 | 39.0 | 49.5 | 38.6 | 40.8 | 38.0 | 51 |
| 52 | 38.0 | 34.5 | 45.0 | 53.0 | 44.2 | 42.4 | 44.1 | 52 |
| 53 | 38.0 | 46.0 | 46.0 | 55.0 | 47.4 | 49.4 | 44.5 | 53 |
| 54 | 38.0 | 47.0 | 56.0 | 59.0 | 47:4 | 50.2 | 44.5 | 54 |
| 55 | 99.9 | 47.0 | 58.0 | 61.5 | 60.4 | 60.8 | 58.8 | 55 |
| .56 | 99.9 | 99.9 | 71.5 | 70.0 | 61.6 | 64.3 | 58.8 | 56 |
| 57 | 99.9 | 99.9 | 72.5 | 71.5 | 61.6 | 64.3 | 99.9 | 57 |
| 58 | 99:9 | 99.9 | 72.5 | 78.0 | 99.9 | 99.9 | 99.9 | 58 |
| 59 | 99.9 | 99.9 | 99.9 | 80.5 | 99.9 | 99.9 | 99.9 | 59 |
| 60 | 99.9 | 99.9 | 99.9 | 80.5 | 99.9 | 99.9 | 99.9 | 60 |
| 61 | 99.9 | 99.9 | 99.9 | 91.5 | 99.9 | 99.9 | 99.9 | 61 |
| 62 | 99.9 | 99.9 | 99.9 | 92.0 | 99.9 | 99.9 | 99.9 | 62 |
| 63 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 63 |

Table C.2. T Scores Obtained by Cumulative Percentages of the Normative Samples: Role Limitations due to Physical Health Problems Scale (RLP)

| T | | Age-Base | d Sample | | Age-S | Stratified S | ample | Т |
|-------|-------|----------|----------|-------|--------------|--------------|--------------|-----------|
| Score | 18-24 | 25-44 | 45-64 | ≥65 | Overall | Female | Male | Score |
| ≤21 | 3.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | |
| 22 | 3.5 | 8.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 <0.1 | ≤21 22 |
| 23 | 3.5 | 8.0 | <0.1 | <0.1 | <0.1 | <0.1 | u- -\0.1 | 22 23 |
| 24 | 7.0 | 8.0 | <0.1 | -<0,1 | ≺ 0:1 | <0.1 | <0.1 | 23 |
| 25 | 7.0 | 8.0 | <0.1 | <0.1 | <0.1 | <0.1 | 9.8 | 25 |
| 26 | 7.0 | 8.0 | <0.1 | <0.1 | 9.4 | 9.0 | 9.8 | 26 |
| 27 | 7.0 | 8.0 | 9.0 | <0.1 | 9.4 | 9.0 | 9.8 | 27 |
| .28 | 7:0 | 8.0 | 9.0 | <0.1 | 9.4 | 9.0 | 9.8 | 28 |
| 29 | 7.0 | 8.0 | 9.0 | <0.1 | 9.4 | 9.0 | 9.8 | 29 |
| 30 | 7:0 | 11.5 | 9.0 | <0.1 | 9:4 | 9.0 | 9.8 | 30 |
| 31 | 7.0 | 11.5 | 9.0 | <0.1 | 9.4 | 9.0 | 9.8 | 31 |
| .32 | 7.0 | 11.5 | 9.0 | <0.1 | 9.4 | 9.0 | 9.8 | 32 |
| 33 | 7.0 | 11.5 | 9.0 | 18.5 | 15.2 | 16.1 | 14.3 | 33 |
| 34 | 12.0 | 11.5 | 18.5 | 18.5 | 15.2 | 16.1 | 14.3 | 34 |
| 35 | 12.0 | 11.5 | 18.5 | 18.5 | 15.2 | 16.1 | 14.3 | 35 |
| 36 | 12.0 | 11.5 | 18.5 | 18.5 | 15.2 | 16,1 | 14.3 | 36 |
| 37 | 12.0 | 11.5 | 18.5 | 18.5 | 15.2 | 16.1 | 14.3 | 37 |
| 38 | 12.0 | 11.5 | 18.5 | 18.5 | 15.2 | 16.1 | 14.3 | 38 |
| 39 | 12.0 | 16.5 | 18.5 | 26.0 | 15.2 | 16.1 | 14.3 | 39 |
| 40 | 12.0 | 16,5 | 18.5 | 26.0 | 15.2 | 16.1 | 20:4 | 40 |
| 41 | 12.0 | 16.5 | 18.5 | 26.0 | 20.2 | 20.0 | 20.4 | 41 |
| 42 | 12:0 | 16.5 | 24:0 | 26.0 | 20,2 | 20.0 | 20.4 | 42 . |
| 43 | 12.0 | 16.5 | 24.0 | 26.0 | 20.2 | 20.0 | 20.4 | 43 |
| 44 | 20.0 | 16.5 | 24.0 | 26:0 | 20.2 | 20:0 | 20.4 | 44 |
| 45 | 20.0 | 16.5 | 24.0 | 34.5 | 20.2 | 20.0 | 20.4 | 45 |
| 46 | 20.0 | 16.5 | 24.0 | 34.5 | 20.2 | 20.0 | 20.4 | 46 |
| 47 | 20.0 | 23.0 | 24.0 | 34.5 | 20.2 | 20.0 | 20.4 | 47 |
| 48 | 20.0 | 23,0 | 24.0 | 34.5 | 29.0 | 30.2 | 27.8 | 48 |
| 49 | 20.0 | 23.0 | 34.0 | 34.5 | 29.0 | 30.2 | 27.8 | 49 |
| 50 | 20.0 | 23.0 | 34.0 | 34.5 | 29.0 | 30.2 | 27.8 | 50 |
| 51 | 20.0 | 23.0 | 34.0 | 34.5 | 29.0 | 30.2 | 27.8 | 51 |
| 52 | 20.0 | 23.0 | 34.0 | 45:0 | 29.0 | 30.2 | 27.8 | 52 |
| 53 | 20.0 | 23.0 | 34.0 | 45.0 | 29.0 | 30.2 | 27.8 | 53 |
| 54 | 99.9 | 23.0 | 34.0 | 45.0 | 29.0 | 30.2 | 27.8 | 54 |
| 55 | 99.9 | 99.9 | 34.0 | 45.0 | 29.0 | 30.2 | 99.9 | 55 |
| 56 | 99.9 | 99.9 | 99.9 | 45.0 | 99.9 | 99.9 | 99.9 | 56 |
| 57 | 99.9 | 99.9 | 99.9 | 45.0 | 99.9 | 99.9 | 99.9 | 57 |
| -58 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 58 |

Table C.3. T Scores Obtained by Cumulative Percentages of the Normative Samples: Pain Scale (PA)

| т | | Age-Base | d Sample | | Age-S | T | | |
|-----------------|-------|----------|----------|--------|---------|--------|------|-------|
| Score | 18-24 | 25–44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| ≤12 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≤12 |
| · 13 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | -<0.1 | <0.1 | 13 |
| 14 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 14 |
| 15 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 15 |
| 16 | 1.0 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 16 |
| 17 | 1.5 | 1.0 | <0.1 | · <0.1 | <0.I | <0.1 | <0.1 | 17 |
| 18 | 1.5 | 1.0 | <0.I | <0.1 | <0.i | <0.1 | 0.4 | 18 |
| 19 | 1.5 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | 0.4 | - 19 |
| 20 | 1.5 | 2.0 | <0.1 | <0.1 | 0.6 | <0.1 | 0.4 | 20 |
| 21 | 1.5 | 2.0 | <0.1 | 1,0> | 0.6 | 8.0 | 0.4 | - 21 |
| 22 | 1.5 | 2.0 | 0.5 | <0.1 | 0.8 | 0.8 | 1.2 | 22 |
| 23 | 3.0 | 4.0 | 0.5 | <0.1 | 1.2 | 1.2 | 1:2 | 23 |
| 24 | 3.0 | 4.0 | 0.5 | <0.1 | 1.2 | 1.2 | 2.9 | 24 |
| 25. | 3.0 | 4.0 | 0.5 | <0.I | 1.2 | 1.2 | 2.9 | 725 |
| 26 | 3.0 | 4.0 | 0.5 | <0.1 | 4.6 | 1.2 | 2.9 | 26 |
| 27 | 3.0 | 4.0 | 5.5 | <0.1 | 4:6 | 6.3 | 3.3 | 27 |
| 28 | 3.5 | 7.0 | 5.5 | 3.0 | 5.0 | 6.3 | 3.3 | 28 |
| 29 | 6.0 | 7.5 | 5.5 | 3.0 | 5.0 | .6.7 | 7.3 | 29 |
| 30 | 6.0 | 7.5 | 6.5 | 3.0 | 7.8 | 6.7 | 7.8 | 30 |
| .31 | 6.0 | 7.5 | 6.5 | 3.0 | 8.0 | 8.2 | 7.8 | 31 |
| 32 | 6.5 | 7.5 | 9.5 | 4.0 | 8.0 | 8.2 | 7.8 | 32 |
| .33 | 6.5 | 8.5 | 9.5 | 8:0 | 8.0 | 8.2 | 7.8 | 33 |
| 34 | 12.0 | 8.5 | 9.5 | 8.0 | 8.0 | 8.2 | 12.2 | 34 |
| 35 | 12.0 | 8.5 | 9.5 | 8.0 | 13.4 | 9:0 | 12.2 | 35 |
| 36 | 12.0 | 8.5 | 9.5 | 8.0 | 13.4 | 14.5 | 12.2 | 36 |
| 37 | 12.0 | 8.5 | 16.0 | 20.0 | 13.4 | 14.5 | 12.2 | 37 |
| 38 | 14.0 | 13.5 | 16.0 | 20.0 | 13.4 | 14.5 | 12.2 | 38 |
| 39 | 17.0 | 13.5 | 16.0 | 20.0 | 13.4 | 14.5 | 18.8 | 39 |
| 40 | 17.0 | 13.5 | 16.0 | 20.0 | 19.0 | 19.2 | 18.8 | 40 |
| 41 | 17.0 | 14.5 | 22.0 | 20.0 | 19.0 | 19.2 | 18.8 | :41 |
| 42 | 17.0 | 25.5 | 22.0 | 28.5 | 19.0 | 19.2 | 19.6 | 42 |
| -4 3 | 24.5 | 25.5 | 22.0 | 28.5 | 21.0 | 22.4 | 28.6 | 43 |
| 44 | 24.5 | 25.5 | 24.5 | 28.5 | 29.8 | 31.0 | 28.6 | 44 |
| 45 | 24.5 | 25.5 | 34.5 | 32.0 | 29.8 | 31.0 | 28.6 | 45 |
| 46 | 24.5 | 29.0 | 34.5 | 41.0 | 29.8 | 31.0 | 28.6 | 46 |
| 47 | 24.5 | 29.0 | 34.5 | 41.0 | 33.6 | .34.9 | 32.2 | 47 |
| 48 | 34.0 | 29.0 | 39.0 | 41.0 | 33.6 | 34.9 | 32.2 | 48 |
| 49 | 34.0 | 29.0 | 39.0 | 45.0 | 33.6 | 34.9 | 32.2 | 49 |
| 50 | 34.0 | 42.0 | 39.0 | 45.0 | 33.6 | 34.9 | 32.2 | 50 |

Table C.3. (Continued)

| T | | Age-Based | d Sample | | Age- | Stratified Sa | ample | T |
|-------|-------|-----------|------------------|------|---------|---------------|-------|---------------|
| Score | 18-24 | 25-44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| SI | 34.0 | 42.0 | 48.5 | 45.0 | 45.6 | 46.7 | 44.5 | ` 5 1. |
| 52 | 60.0 | 42.0 | 48.5 | 45.0 | 45.6 | 46.7 | 44.5 | 52 |
| :53 | 60.0 | 69.0 | 4 8.5 | 56.0 | 45.6 | 46.7 | 44.5 | 53 |
| 54 | 60.0 | 69.0 | 67.0 | 56.0 | 69.2 | 68.2 | 70.2 | 54 |
| 55 | 60.5 | 69.0 | 67.0 | 56.0 | 69.2 | 68.2 | 70.2 | 55 |
| 56 | 60.5 | 69.0 | 67.0 | 77.5 | 69.2 | 68.2 | 70.2 | 56 |
| .57 | 60.5 | 69.0 | 67.0 | 77.5 | 69.2 | 68.2 | 70,2 | 57 |
| 58 | 60.5 | 69.0 | 67.0 | 77.5 | 69.2 | 68.2 | 70.2 | 58 |
| 59 | 99.9 | 99.9 | 67.0 | 78.5 | 69.2 | 68.2 | 70.2 | 59 |
| 60 | 99.9 | 99.9 | 99.9 | 78.5 | 99.9 | 99.9 | 99.9 | 60 |
| 61 | 99.9 | 99.9 | 99.9 | 78.5 | 99.9 | . 99.9 | 99.9 | 61 |
| 62 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 62 |

Table C.4. T Scores Obtained by Cumulative Percentages of the Normative Samples: General Health Perceptions Scale (GHP)

| т | | Age-Based | ł Sample | | Age-S | Stratified Sa | ample | T |
|-------|-------|-----------|----------|------|---------|---------------|-------------|-----------|
| Score | 18–24 | 25-44 | 4564 | ≥65 | Overall | Female | Male | Score |
| ≤17 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≤17 |
| 18 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 18 |
| 19 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 19 |
| 20 | 0:5 | 0.5 | <0.1 | <0.1 | <0.1 | 0.4 | <0.1 | 20 |
| 21 | 0.5 | 1.0 | <0.1 | <0.1 | 0.2 | 0.4 | <0.1 | 21 |
| 22 | 0.5 | 1.0 | 0.5 | <0.1 | 0.2 | .0.4 | <0.1 | 22 |
| 23 | 1.0 | 2.5 | 0.5 | <0.1 | 0.4 | 2.0 | <0.1 | 23 |
| 24 | - 1.5 | 2.5 | 0.5 | <0.1 | 0.8 | 2.0 | 0.4 | 24 |
| 25 | 1.5 | 2.5 | 0.5 | <0.1 | 1.6 | 2.4 | 0.4 | 25 |
| 26 | 3,0 | 3.0 | 0.5 | 0.5 | 1.8 | 3.1 | 1.2 | 26 |
| 27 | 3.0 | 4.0 | 1.5 | 0.5 | 2.6 | 3.5 | 1.2 | 27 |
| 28 | 3.5 | ·4.0 | 1.5 | 1.0 | 3.8 | 3.5 | 1.6 | 28 |
| 29 | 4.0 | 4.0 | 2.5 | 1.5 | 4.0 | 3.5 | 4.5 | 29 |
| .30 | 4.0 | 5.5 | 4.5 | 2.5 | 4.4 | 4.3 | 4.5 | 30 |
| 31 | 6.0 | 5.5 | 5.0 | 2.5 | 6.0 | 5.1 | 5.3 | 31 |
| . 32 | 6.5 | 6:0 | 7.0 | 3.5 | 6.8 | 6.7 | ~6.5 | _32 |
| 33 | 7.0 | 7.0 | 8.0 | 4.5 | 7.8 | 8.2 | 8.6 | 33 |
| 34 | 8.0 | 7.5 | 9.5 | 6.5 | 9.2 | 9.0 | 9.0 | 34 |
| 35 | 8.0 | 8.5 | 12.0 | 6.5 | 10.4 | 10.2 | 10.2 | 35 |
| 36 | 9.5 | 10.0 | 13.5 | 9.5 | 11.4 | 11.0 | 11.8 | 36 |
| 37 | 11.5 | 11.5 | 13.5 | 13.5 | 12.6 | 12.2 | 12.7 | 37 |
| .38 | 13.5 | 12.5 | 15.5 | 16.0 | 13.8 | 13.7 | 13.9 | 38 |
| 39 | 16.0 | 15.0 | 17.0 | 19.5 | 15.4 | 14.9 | 15.5 | 39 |
| 40 | 17.0 | 17.0 | 18.0 | 22.0 | 17.6 | 16.5 | 18.0 | 40 |
| 41 | 20.5 | 18.5 | . 20.5 | 24.0 | 18.4 | 18.0 | 19.6 | 41 |
| 42 | 23.0 | 22.0 | 22.5 | 24.5 | 20.2 | 19.6 | 20.0 | 42 |
| 43 | 24.0 | 24.5 | 24.0 | 26.0 | 23.2 | 22.4 | 22.9 | 43 |
| 44 | 27:0 | .28.5 | 26.0 | 29.0 | 25.6 | 25.9 | 27.3 | 44 |
| 45 | 30.0 | 29.0 | 27.5 | 31.0 | 29.2 | 30.6 | 31.0 | 45 |
| 46 | 36.0 | 34.5 | 33,5 | 36.0 | 33.6 | 31.8 | 36.3 | 46 |
| 47 | 37.0 | 35.5 | 40.0 | 40.0 | 36.8 | 35.7 | 37.6 | 47 40 |
| 48 | 41.0 | 38.0 | -41.5 | 45.5 | 41.0 | 38.0 | 42.0 | 48 |
| 49 | 41.5 | 40.0 | 45.0 | 46.5 | 43.0 | 42.7 | 46.5 | 49 50 |
| 50 | 49.0 | 45.5 | 48.5 | 49.5 | 47.2 | 44.3 | 50.2 | 50 51 |
| 51 | 52.0 | 48.5 | 53.5 | 54.5 | 51.8 | 51.4 | 51.0 | 51 52 |
| 52 | 56.0 | 51.0 | 54.0 | 58.0 | .55.2 | 53.7 | 57.1 | 52 |
| 53 | 56.5 | 62.0 | 60.0 | 61.5 | 57.8 | 60.0 | 58.8 | 53 |
| 54 | 64:5 | 66.5 | 62.0 | 67.5 | 66.2 | 65.5 | 65.7 | 54 |
| 55 | 66.5 | 67.5 | 70.5 | 69.5 | 67.2 | 71.4 | 68.2 | 55 ••• |
| 56 | 68.5 | 70.5 | 71.5 | 72.0 | 72.4 | 72.2 | 70.6 | 56 |

Table C.4. (Continued)

| T | | Age-Base | d Sample | | Age- | Stratified S | ample | T |
|-------|-------|----------|----------|------|---------|--------------|-------|-------|
| Score | 18–24 | 25-44 | 4564 | ≥65 | Overall | Female | Male | Score |
| 57 | 74.5 | 74.0 | 73.5 | 76.5 | 74.4 | 73.3 | 75.5 | 57 |
| 58 | 77.0 | 78.5 | 75.0 | 77.5 | 78.4 | 80.4 | 77.6 | -58 |
| 59 | 79.0 | 78.5 | 81.0 | 82.0 | 79.6 | 82.7 | 78.8 | 59 |
| 60 | 81.5 | 82.5 | 81.0 | 83.5 | 82.0 | 83.1 | 80.8 | 60 |
| 61 | 84.0 | 85.5 | 84.5 | 84.5 | 85.6 | 86.3 | 84.9 | 61 |
| 62 | 84.0 | 85.5 | 89.0 | 87.0 | 87.8 | 88.2 | 87.3 | 62 |
| 63 | 99.9 | 99.9 | 91.5 | 89.0 | 87.8 | 88.2 | 87.3 | 63 |
| 64 | 99.9 | 99.9 | 91.5 | 90.0 | 99.9 | 99,9 | 99.9 | 64 |
| 65 | 99.9 | 99.9 | 99.9 | 93.0 | 99.9 | 99.9 | 99.9 | 65 |
| 66 | 99.9 | 99.9 | 99.9 | 94.5 | 99.9 | 99:9 | 99.9 | 66 |
| 67 | 99.9 | 99.9 | 99.9 | 94.5 | 99.9 | 99.9 | 99.9 | 67 |
| -68 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99:9 | 99.9 | 68 |

Table C.5. T Scores Obtained by Cumulative Percentages of the Normative Samples: Emotional Well-Being Scale (EWB)

| T | | Age-Based | l Sample | | Age-S | · T | | |
|----------|-------|---------------|----------|-------|---------|--------------|------|-------|
| Score | 18-24 | 25 <u>–44</u> | 45–64 | ≥65 | Overall | Female | Male | Score |
| ≤16 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≤16 |
| 17 | <0.1 | <0.1 | <0.1 | <0.1 | 0.2 | ≺ 0.1 | 0.4 | 17 |
| 18 | <0.1 | <0.1 | <0.1 | <0.1 | 0.4 | <0.1 | 0.8 | 18 |
| 19 | <0.1 | <0.1 | <0.1 | <0.1 | 0.6 | 0.4 | 0.8 | 19 |
| 20 | 0.5 | <0.1 | 0.5 | 0.5 | 0.6 | 0.4 | 0.8 | 20 |
| 21 | 0.5 | 0.5 | 0.5 | 0.5 | 0.8 | 0.4 | 0.8 | 21 |
| 22 | 0.5 | 1.0 | 0.5 | 0.5 | 1.0 | 0.8 | 8.0 | 22 |
| 23 | 0.5 | 1.0 | 0.5 | 1.0 | 1.6 | 1.2 | 0.8 | 23 |
| 24 | 1.5 | 1.0 | 1.0 | 1.0 | 1.8 | 2.7 | 1.2 | 24 |
| .25 | ±1.5 | 1.5 | 2.0 | 1.0 | 2.2 | 3.1 | 2.0 | .25 |
| 26 | 2.0 | 2.5 | 2.0 | 2.0 | 2.8 | 3.1 | 2,4 | 26 |
| 27 | 2.0 | 3.0 | 2.5 | 2.5 | 3.0 | 3.5 | 2.4 | 27 |
| 28 | 2.5 | 4.0 | 3.5 | 3.0 | 3.4 | 3.9 | 2.4 | 28 |
| 29 | 3.5 | 4.5 | 4.5 | 3.5 | 3.6 | 4.3 | 3.7 | 29 |
| 30 | 4.0 | 5.0 | 6.0 | 5.0 | 4.4 | 4.7 | 3.7 | 30 |
| 31 | 5.0 | 5.0 | 7,5 | 5,0 | 5.0 | 5.9 | 5.7 | 31 |
| 32 | 5.5 | 6.0 | 8.5 | 6.5 | 6.6 | 6.7 | 6.1 | 32 |
| 33 | 7.0 | 6.5 | 9.5 | 8.0 | 7.6 | 7.5 | 6.5 | 33 |
| 34 | 7.5 | 7.5 | 10.5 | 8.5 | 8.4 | 9.4 | 7.3 | 34 |
| .35 | 8.5 | 9.0 | 10.5 | 11.0 | 8.6 | 9.4 | 10.2 | 35 |
| 36 | 10.0 | 11.5 | 10.5 | 12.5 | 10.8 | 10.2 | 11.8 | 36 |
| 37 | 11.0 | 12.5 | 13.0 | 14.0 | 12:2 | 11.0 | 14.3 | 37 |
| 38 | 14.0 | 15.0 | 14.0 | 14.0 | 14.6 | 11.4 | 15.1 | 38 |
| 39 | 15.5 | 16.5 | 16.5 | 16.0 | 15.6 | 15.3 | 18.0 | 39 |
| 40 | 18.5 | 18.0 | 19.5 | 18.5 | 19.4 | 16.1 | 20.0 | 40 |
| 41 | 20,0 | 19.5 | 21.0 | 20.0 | 20:4 | 20.0 | 20.4 | 41 |
| 42 | 23.0 | 22.0 | 23.5 | 22.5 | 22.4 | 22.0 | 21.6 | 42 |
| 43 | 25.5 | 25.0 | 24.0 | 23.5 | 23.8 | 23.5 | 23.7 | 43 |
| 44 | 28.0 | 27.5 | 28.0 | 26.5 | 27.0 | 26.7 | 25.7 | 44 |
| 45 | 30.5 | 28.5 | 28.5 | 27.5 | .28.4 | 29.8 | 28.6 | 45 |
| 46 | 34.0 | 33.5 | 31.0 | 32.0 | 32.0 | 31.8 | 29.0 | 46 |
| -47 | 35.5 | 34.5 | 32.0 | 33.0 | 32.6 | .34.9 | 35.1 | 47 |
| 48 | 41.0 | 38.5 | 35.0 | 38.5 | 37.8 | 38.4 | 36.3 | 48 |
| 49 | 42.0 | 44.5 | 41.0 | 42.5 | 40.8 | 40.4 | 40.8 | 49 |
| 50 | 49.5 | 47.0 | 43.0 | 48.0 | 44.8 | 44.3 | 43.3 | 50 |
| .5.1 | 51:0 | 49.5 | 44.5 | .51.0 | 46.2 | 47.8 | 45.3 | 51 |
| 52 | 56.5 | 53.5 | 47.5 | 51.0 | 51.6 | 50.6 | 48.6 | 52 |
| 53 | 57.5 | 55.0 | 55.0 | 60.5 | 52.8 | 55.7 | 57.6 | 53 |
| 54 | 62.0 | 60.0 | 58.0 | 60.5 | 62.4 | 60.8 | 60.8 | 54 |

Table C.5. (Continued)

| T | | Age-Based | d Samula | | | | | |
|----------|-------|-----------|----------|--|---------|----------------------------|--|------------------------|
| Score | 10.24 | | <u> </u> | | Age-S | tratified Sa | ample | T |
| | 18–24 | 25-44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| 55 | 63.0 | 63.5 | 70.0 | 62.0 | 62.8 | 64.7 | 67.3 | 55 |
| 56 | 74.0 | 68.0 | 70.0 | 77.0 | 75.2 | 71.4 | 72.2 | 56 |
| 57 | 76.5 | 74.0 | 83,0 | 78.5 | 75.2 | 78.0 | 73.9 | :57 |
| 58 | 85.0 | 80.5 | 83.5 | 80.0 | 86.8 | 78.4 | 84.5 | 58 |
| 59 | 85.0 | 81.5 | 85.5 | 80.0 | 87.0 | 89.4 | 85.3 | 59 |
| 60 | 86.0 | 90.0 | 85.5 | 88.5 | 89.0 | 89.4 | 87.3 | 60 |
| 61. | 90.0 | 91.0 | 92.5 | 88.5 | 89,0 | 90:6 | 87.8 | BROSERS SPANISHED LAST |
| 62 | 92.5 | 91.0 | 93.0 | 89.5 | 94.6 | 91.0 | 93.9 | 61 |
| 63 | 92.5 | 96:5 | 93.5 | 89.5 | 94,6 | 95.3 | 94.3 | 62 |
| 64 | 92.5 | 96.5 | 93.5 | 89.5 | 95.6 | 96.1 | 94.7 | 63 |
| 65 | 96.0 | 97.5 | 99.9 | 99.9 | 95.6 | 96.5 | September of the septem | 64 |
| 66 | 96.0 | 97.5 | 99.9 | 99.9 | 99.9 | 96.5 | 94.7 | 65 |
| -67 | 96.0 | 99.9 | 99.9 | 99.9 | 99.9 | Chia Santa de Maria (1940) | 99.9 | 66 |
| 68 | 96.0 | 99.9 | 99.9 | 99.9 | | 99.9 | 99:9 | 67 |
| 69 | 99.9 | 99.9 | 99.9 | NO SECTION AND ADDRESS OF THE PARTY OF THE P | 99.9 | 99.9 | 99.9 | 68 |
| | | 1.5.7 | 77,7 | 99.9 | 99.9 | 99.9 | 99.9 | 69 |

Table C.6. T Scores Obtained by Cumulative Percentages of the Normative Samples: Role Limitations due to Emotional Problems Scale (RLE)

| T | | Age-Base | d Sample | | <0.1 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 5.7 6.4 <0.1 10.6 11.6 12.5 10.6 11.6 12.5 10.6 11.6 12.5 10.6 11.6 12.5 10.6 11.6 12.5 10.6 11.6 12.5 10.6 11.6 12.5 10.6 11.6 12.5 10.6 11.6 12.5 10.6 | | | т |
|-------------|-------|----------|----------|-------|--|---|-------------------------------|------------|
| Score | 18–24 | 25-44 | 45-64 | ≥65 | | | | Score |
| ≤I8 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 5.7 | ≤18 |
| 19 | 5.5 | <0.1 | 7.0 | <0.1 | 6.4 | <0.1 | 81%S675456G000100000000000000 | 19 |
| 20 | 5.5 | <0.1 | 7.0 | <0.1 | 6.4 | <0.1 | 5.7 | 20 |
| .21 | 5.5 | <0.1 | 7.0 | <0.1 | 6.4 | 7,] | 5.7 | 21 |
| 22 | 5.5 | <0.1 | 7.0 | <0.1 | 6.4 | 7.1 | 5.7 | 22 |
| 23 | .5.5 | <0,1 | 7.0 | <0.1 | 6:4 | 7.1 | 5.7 | 23 |
| 24 | 5.5 | 8.5 | 7.0 | <0.1 | 6.4 | 7 .1 | 5.7 | 24 |
| 25 | 5,5 | 8.5 | 7.0 | 11.0 | 6.4 | 7.1 | 5.7 | 25 |
| 26 | 5.5 | 8.5 | 7.0 | 11.0 | 6.4 | 7.1 | 5.7 | 26 |
| .27 | 5.5 | 8.5 | 7.0 | 11.0 | 6.4 | 7.1 | 5.7 | 27 |
| 28 | 5.5 | 8.5 | 7.0 | 11.0 | 6.4 | 7.1 | 5.7 | 28 |
| 29 | 5.5 | 8.5 | 7,0 | 0.11 | 6,4 | 7.1 | 10.6 | 29 |
| 30 | 5.5 | 8.5 | 7.0 | 11.0 | 6.4 | 7.1 | 10.6 | 30 |
| 31 | 12.0 | 8.5 | 11.5 | 11.0 | 11.6 | 7.1 | 10.6 | 31 |
| 32 | 12.0 | 8.5 | 11.5 | 11.0 | 11.6 | 12.5 | 10.6 | 32 |
| -33 | 12.0 | 8.5 | 11.5 | 11.0 | 11.6 | 12.5 | 10.6 | 33 |
| 34 | 12.0 | 16.5 | 11.5 | 11.0 | 11.6 | 12.5 | 10.6 | 34 |
| .35 | 12.0 | 16.5 | 1.1.5 | 15:0 | 11.6 | 12.5 | 10.6 | 35 |
| 36 | 12.0 | 16.5 | 11.5 | 15.0 | 11.6 | 12.5 | 10.6 | 36 |
| .37 | 12.0 | 16.5 | 11.5 | 15.0 | 11.6 | 12.5 | 10.6 | .37 |
| 38 | 12.0 | 16.5 | 11.5 | 15.0 | 11.6 | 12.5 | 10.6 | 38 |
| 39 | 12.0 | 16.5 | 11.5 | 15.0 | 11.6 | 12.5 | 10.6 | .39 |
| 40 | 12.0 | 16.5 | 11.5 | 15.0 | POSE SHIP SHIP SHIP SHIP SHIP SHIP SHIP SHIP | 12.5 | 10.6 | 40 |
| 41 | 12.0 | 16.5 | 41.5 | 15.0 | 11.6 | 12.5 | .10.6 | 41 |
| 42 | 12.0 | 16.5 | 11.5 | 15.0 | MARKET STATE OF THE STATE OF TH | 0.0000000000000000000000000000000000000 | 900968/00G088860B:0508960C | 4 2 |
| -:43 | 22.5 | 16.5 | 17.0 | 15.0. | R 026 904 042 050 000 000 000 | 22.4 | 15.5 | 43 |
| 44 | 22.5 | 16.5 | 17.0 | 15.0 | Stort in a property of the store of the stor | | 95450501 WASANGANGANA | 44 |
| 45 | 22.5 | 24.0 | 17.0 | 22.5 | 19:0 | 22.4 | 15.5 | 45 |
| 46 | 22.5 | 24.0 | 17.0 | 22.5 | SECONO CON CONTRACTOR CONTRACTOR | 0.77353.07002.085394.053 | 15.5 | 46 |
| 47 | 22.5 | .24:0 | 17.0 | 22.5 | | 16079A00959A06503H6215S16 | | -47 |
| 48 | 22.5 | 24.0 | 17.0 | 22.5 | Programme and the second | y consultation of the consultation of the | 15.5 | 48 |
| .49 | 22.5 | 24:0 | 17.0 | 22.5 | | | | -49 |
| 50 | 22.5 | 24.0 | 17.0 | 22.5 | 19.0 | 22.4 | 15.5 | 50 |
| 51 | 22.5 | 24.0 | 17.0 | 22.5 | 19.0 | 22:4 | 15.5 | .51 |
| 52 | 22.5 | 24.0 | 17.0 | 22.5 | 19.0 | 22.4 | 15.5 | 52 |
| 53 | 22.5 | 24.0 | 17.0 | 22.5 | 19.0 | 22.4 | 15.5 | 53 |
| 54 | 22.5 | 24.0 | 99.9 | 22.5 | 99.9 | 22.4 | 99.9 | 54 |
| .55 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | .55 |

Table C.7. T Scores Obtained by Cumulative Percentages of the Normative Samples: Social Functioning Scale (SF)

| T | | Age-Based | d Sample | | Age- | Stratified S | ample | т |
|------------|-------|-----------|----------|-----------------|---------|--------------|--------------------|-----------------------|
| Score | 18-24 | 25-44 | 45-64 | ≥65 | Overall | Female | Male | Score |
| 7 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 7 |
| 8 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | .8 |
| 9 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 9 |
| 10 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 10 |
| | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | 0.4 | <0.1 | 11 |
| 12 | 0.5 | <0.1 | <0.1 | <0.1 | 0.2 | 0.4 | <0.1 | 12 |
| 13 | 0.5 | <0.1 | <0.1 | <0.1 | 0.2 | 0.4 | <0.1 | 13 |
| 14 | 0.5 | <0.1 | ·<0.1 | <0.1 | 0.2 | 0.4 | . <0;1 | 14 |
| 15 | 0.5 | 0.5 | <0.1 | <0.1 | 0.4 | 0.4 | <0.1 | 15 |
| 16 | 0.5 | 0.5 | 0,5 | <0.1 | 0,4 | 0.4 | 0:4 | 16 |
| 17 | 1.0 | 0.5 | 0.5 | <0.1 | 0.8 | 0.8 | 0.4 | 17 |
| 18 | 1.0 | 0.5 | 1.0 | < 0.1 | 0.8 | .0.8 | 0.8 | 18 |
| 19 | 1.0 | 0.5 | 1.0 | <0.1 | 1.0 | 0.8 | 0.8 | 19 |
| 20 | 1:0 | .0.5 | 2.0 | <0.1 | 2.0 | 1.6 | 1.2 | 20 |
| 21 | 1.0 | 0.5 | 2.0 | 2.0 | 2.0 | 1.6 | 2.4 | 21 |
| 22 | 1:0 | 1.0 | 2.5 | 2.0 | 2.0 | 1.6 | 2.4 | 21 |
| 23 | 1.0 | 3.5 | 4.0 | 2.0 | 2.0 | 1.6 | 2.4 | 23 |
| 24 | 1.5 | 3.5 | 4.0 | 4.0 | 4.2 | 3.1 | 2.4 | 23 |
| 25 | 2.0 | 3.5 | 4.0 | 4.0 | 4.2 | 3.1 | 5.3 | 2 5 |
| 26 | 2.0 | 3.5 | 4.0 | 4.0 | 4.2 | 3.1 | 5.3 | 26 |
| 27 | 2.0 | 6.5 | 6.5 | 4.0 | 5.4 | 4.3 | 5.3 | 27 |
| 28 | 2.0 | 6.5 | 6.5 | 4.0 | 5.4 | 4.3 | 6.5 | CONTRACTOR CONTRACTOR |
| 29 | 2.0 | 6.5 | 6.5 | 9.0 | 5.4 | 4.3 | 6.5 | 28 29 |
| 30 | 7.0 | 7.5 | 7.0 | 9.0 | 5.4 | 4.3 | 6.5 | 30 |
| 31 | 7.0 | 7.5 | 7.0 | 9.0 | 9.6 | 10.2 | 6.5 | 31 |
| 32 | 7.0 | 7.5 | 7.0 | 9.5 | 9.6 | 10:2 | 9.0 | 32 |
| 33 | 7.5 | 10.5 | 12.0 | 9.5 | 9.6 | 10.2 | 9.0 | 33 |
| 34 | 7.5 | 10.5 | 12.0 | 10.0 | 10.4 | 11.4 | 9.0 | 34 |
| 35 | 12.5 | 10.5 | 12.0 | 11.0 | 10.4 | 11.4 | 9.4 | 35 |
| 36 | 13.5 | 11.0 | 12.5 | 11.0 | 13.6 | 14.1 | 9.4 | 36 |
| 37 | 21.5 | 11.0 | 12.5 | 16.0 | 13.8 | 14.1 | 13.5 | 37 |
| 38 | 21,5 | 16.0 | 13.5 | 17.0 | 13.8 | 18.8 | 13.5 | 38 |
| 39 | 21.5 | 16.0 | 13.5 | 17.0 | 19.8 | 20.0 | 19.2 | 39 |
| 40 | 21:5 | 21.5 | 19.5 | 17.0 | 19.8 | 20.0 | 19.6 | 40 |
| 41 | 21.5 | 22.5 | 19.5 | 17.0 | 19.8 | 20.0 | 19.6 | 41 |
| 4 2 | 21:5 | 22.5 | 19.5 | 20.0 | 19.8 | 20.0 | 19.6 | CASCADIA PROGRAMMA |
| 43 | 23.0 | 22.5 | 19.5 | 22.0 | 19.8 | 20.0 | 19.6 | 42 |
| 44 | 23.5 | 22.5 | 19.5 | 22,0 | 22.8 | 22.7 | 22.9 | 43 44 |
| 4 5 | 30.5 | 24.0 | 24.5 | 28.0 | 22.8 | 22.7 | PART AUGUST STATES | 44 |
| 46 | 30.5 | 24.0 | 24.5 | 28.0 | 29.8 | 29.8 | 22.9 29.8 | 45 46 |

Table C.7. (Continued)

| T | | Age-Base | d Sample | | Age-S | Stratified Sa | ample | т |
|----------|---------|----------|----------|------|---------|---------------|-------|-------|
| Score | 18–24 | 25-44 | 4564 | ≥65 | Overall | Female | Male | Score |
| 47 | 30.5 | 33.5 | 30.0 | 28.0 | 29.8 | 29.8 | 29.8 | 47 |
| 48 | 30.5 | 33.5 | 30.0 | 28.0 | 29.8 | 29.8 | 29.8 | -48 |
| 49 | 30.5 | 33.5 | 30.0 | 28.0 | 29.8 | 29.8 | 29.8 | 49 |
| 50 | 30.5 | 33.5 | 30.0 | 35.5 | 29.8 | 29.8 | 29.8 | 50 |
| 51 | 41.5 | 33.5 | 37.0 | 39.0 | 36.0 | 38.0 | 37.6 | 51 |
| 52 | ÷41.5 ÷ | 47.0 | 41.5 | 39.0 | 40.8 | 43.9 | 37.6 | 52 |
| 53 | 41.5 | 47.0 | 41.5 | 39.0 | 40.8 | 43.9 | 37.6 | 53 |
| 54 | 41.5 | 47.0 | 4L5 | 39.0 | ÷40.8 | 43,9 | 37.6 | 54 |
| 55 | 41.5 | 47.0 | 41.5 | 39.0 | 40.8 | 43.9 | 37.6 | 55 |
| 56 | 41.5 | '47.0 | 99.9 | 99.9 | 40.8 | 43.9 | 99.9 | 56 |
| 57 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 57 |

Table C.8. T Scores Obtained by Cumulative Percentages of the Normative Samples: Energy/Fatigue Scale (EF)

| T | | | d Sample | | Age- | Stratified : | Sample | т |
|-------|-------|-------|----------|------|---------|--------------|--------|-------|
| Score | 18–24 | 25-44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| 25 | <0.1 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | 1.2 | 25 |
| 26 | 1.5 | 0.5 | <0.1 | 1.5 | 1.4 | 1.6 | 1.2 | 26 |
| 27 | 1.5 | 0.5 | 1.0 | 1.5 | 1.4 | 1.6 | 1.6 | 27 |
| 28 | 2:0 | 1.0 | 1.0 | 1.5 | 1.6 | 1.6 | 2.4 | 2.8 |
| 29 | 2.5 | 1.5 | 1.0 | 1.5 | 2.2 | 1.6 | 3.7 | 29 |
| 30 | 3.5 | 3.5 | 2.0 | 2.0 | 3.6 | 2.0 | 3.7 | 30 |
| 31 | 4.0 | 5.0 | 3.0 | 5.5 | 5.0 | 3.5 | 4.9 | 31 |
| 32 | 4.0 | .5.0 | 5.0 | 5.5 | 5:4 | 5.9 | 6.1 | 32 |
| 33 | 6.5 | 6.0 | 6.5 | 8.5 | 7.6 | 6.3 | 6.1 | 33 |
| .34 | 7.5 | 6.5 | 8.5 | 9.0 | 8.0 | 9.4 | 7.3 | 34 |
| 35 | 8.5 | 9.0 | 8.5 | 10.0 | 10.0 | 10.6 | 9.0 | 35 |
| .36 | 9.5 | 11.0 | 10.0 | 10.5 | 10:6 | 11.8 | 12.2 | 36 |
| 37 | 11.5 | 15.0 | 12.0 | 11.0 | 13.2 | 12.2 | 13.1 | 37 |
| 38 | 13.5 | 16.5 | 14.0 | 15.0 | 14.4 | 14.5 | 16.3 | 38 |
| 39 | 16.0 | 17.5 | 18.0 | 15.0 | 16.8 | 16.1 | 16.7 | 39 |
| 40 | 17.0 | 18.0 | 19.5 | 18.5 | 18.6 | 17.3 | 18.4 | 40 |
| 41 | 20.0 | 21.5 | 22.5 | 18.5 | 20.4 | 20.8 | 22.9 | 41 |
| 42 | 22.5 | 22.5 | 23.5 | 25.0 | 24.8 | 22.4 | 24.9 | 42 |
| 43 | 27.0 | 27.5 | 30.0 | 25.0 | 27.4 | 27.1 | 26.1 | 43 |
| 44 | 30.5 | 29.5 | 32.0 | 30:5 | 29.6 | 29.8 | 29.0 | 44 |
| 45 | 33.0 | 32.0 | 36.0 | 31.0 | 32.0 | 33.3 | 32.2 | 45 |
| 46 | 38.0 | 34.5 | 39.5 | 38.0 | 36.6 | 34.9 | 34.7 | 46 |
| 47 | 41.5 | 39.0 | 42.5 | 39.0 | 40.0 | 41.2 | 35.5 | 47 |
| ·48 | 43.5 | 41.5 | 43.0 | 43.0 | 40.4 | 45.1 | 42.4 | 48 |
| 49 | 48.0 | 43.5 | 49.0 | 44.5 | 47.4 | 45.1 | 43.7 | 49 |
| .50 | 50.5 | 48.5 | 49.0 | 50.5 | 48.8 | 52.5 | 46.9 | 50 |
| 51 | 53.5 | 51.5 | 50.0 | 51.5 | 52.6 | 53.7 | 48.6 | 51 |
| 52 | 56.0 | 55.5 | 53.5 | 56.0 | 53.8 | 58.0 | 56.3 | 52 |
| 53 | 64.5 | 57.0 | 59.0 | 58.0 | 60.6 | 58.8 | 56.7 | 53 |
| .54 | 66.0 | 65.5 | 59.5 | 61.0 | 61.2 | 64.7 | 59.2 | 54 |
| 55 | 68.0 | 66.5 | 62.0 | 69.0 | 64.0 | 65.5 | 65.7 | 55 |
| 56 | 72.5 | 70.0 | 69.0 | 70.5 | 69.8 | 68.6 | 69.0 | 56 |
| 57 | 74.0 | 75.5 | 71.5 | 72.0 | 72.2 | 73.7 | 71.8 | 57 |
| 58 | 75.0 | 77.0 | 73.5 | 73.0 | 74.4 | 75.3 | 72.2 | 58 |
| 59 | 77.5 | 79.0 | 84.0 | 82.5 | 75.2 | 76.9 | 82.9 | 59 |
| 60 | 87.0 | 79.5 | 84.0 | 83.5 | 85.2 | 78.0 | 83.3 | 60 |
| 61 | 87.5 | 89.0 | 92.0 | 90.0 | 85.6 | 87.5 | 91.8 | 61 |
| 62 | 92.0 | 89.0 | 92.0 | 90.0 | 93.6 | 87.8 | 91.8 | 62 |
| 63 | 92.5 | 94.5 | 94.5 | 94.5 | 94.0 | 95.3 | 96.3 | 63 |

Table C.8. (Continued)

| Τ | | Age-Based | i Sample | | Age-S | Stratified Sa | ample | Τ |
|-------|-------|-----------|--------------|------|---------|---------------|-------|-------|
| Score | 18–24 | 25–44 | 45-64 | ≥65 | Overall | Female | Male | Score |
| 64 | 93.0 | 96.0 | 94.5 | 94.5 | .97.4 | 95.7 | 96.7 | 64 |
| 65 | 95.5 | 96.0 | 94.5 | 94.5 | 97.6 | 98.4 | 96.7 | 65 |
| 66 | 96.5 | 98.5 | 9 6.5 | 96,5 | 97.6 | 98.4 | 98.0 | -66 |
| 67 | 96.5 | 98.5 | 96.5 | 96.5 | 98.2 | 98.4 | 98.4 | 67 |
| 68 | 98.0 | 98.5 | 96.5 | 96.5 | 98.4 | 98.4 | 98.4 | - 68 |
| 69 | 98.0 | 99.0 | 99.9 | 99.9 | 98.4 | 98.4 | 99.9 | 69 |
| 70 | 99,9 | 99.0 | 99.9 | 99.9 | 99.9 | 98,4 | 99.9 | 70 |
| 71 | 99.9 | 99.0 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | ·71 |
| 72 | 99.9 | 99.9 | 99.9 | 99,9 | 99.9 | 99.9 | 99.9 | 72 |

Table C.9. T Scores Obtained by Cumulative Percentages of the Normative Samples: Physical Health Composite (PHC)

| T | | Age-Base | d Sample | | Age-Stratified Sample | | | |
|-----------|-------|----------|----------|------|-----------------------|--------|--------------|---|
| Score | 18-24 | 25–44 | 45–64 | ≥65 | Overali | Female | Male | T Scor |
| ≤7 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≤7 |
| 8 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≤ 0.1 | 8 |
| 9 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 9 |
| 10 | 0.5 | -<0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 10 |
| 11 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 11 |
| 12 | .0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 12 |
| 13 | 0.5 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 13 |
| 14 | 1.5 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | -<0.1 | 14 |
| 15 | 1.5 | 1.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 15 |
| 16 | 1.5 | 1.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 16 |
| 17 | 1.5 | 2.0 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 17 |
| 18 | 1.5 | 2.5 | <0.1 | <0.1 | 0.2 | 0.4 | <0.1 | 18 |
| 19 | 1.5 | 3.0 | <0.1 | <0.1 | 0.6 | 1.2 | 0.4 | 19 |
| 20 | 1.5 | 3.5 | <0.1 | <0.1 | 1.0 | 1.2 | 0.8 | 20 |
| 21 | 1.5 | 4.0 | <0.1 | <0.1 | 1.0 | 1.2 | 0.8 | 21 |
| 22 | 1.5 | 4.5 | <0.1 | <0.1 | 1.4 | 1.6 | 1.2 | 22 |
| 23 | 2.5 | 4.5 | 1.5 | 0.5 | 1.8 | 2.4 | 1.6 | 23 |
| 24 | 3.0 | 4.5 | 1.5 | 0.5 | 3.6 | 4.3 | 2.9 | 24 |
| 25 | 3.5 | 5.0 | 2.0 | 0.5 | 4.4 | 5.1 | 4.1 | 25 |
| 26 | 4.0 | .5.5 | 2.5 | 0:5 | 5.0 | 5.1 | 6.1 | 26 |
| 27 | 5.0 | 6.5 | 4.0 | 0.5 | 5.6 | 5.1 | 6.1 | 27 |
| 28 | 5.5 | 6.5 | 6.0 | 1.5 | 6.0 | 5.1 | 6.9 | 28 |
| 29 | 5.5 | 6.5 | 6.0 | 2.5 | 6.6 | 5.9 | 7.3 | 29 |
| 30 | 6.0 | 6.5 | 7:0 | 5.0 | 7.0 | 6.7 | 8.2 | 30 |
| 31 | 7.0 | 6.5 | 7.0 | 6.0 | 8.4 | 7.5 | 9.4 | 31 |
| .32 | 8.0 | 6.5 | 8.5 | 8.5 | 9.4 | 9.0 | 9.8 | 32 |
| 33 | 8.5 | 7.0 | 0.01 | 9.5 | 10.0 | 9.8 | 10.6 | 33 |
| 34 | 10.5 | 7.0 | 10.5 | 10.5 | 11.2 | 11.0 | 10.6 | 34 |
| 35 | 0.11 | 8.5 | 11.5 | 13.5 | 11.8 | 11.8 | 11.4 | 35 |
| 36 | 12:0 | 9.5 | 13.0 | 15.5 | 12.2 | 12.2 | 11.8 | 36 |
| 37 | 13.0 | 10.5 | 15.5 | 18.0 | 13.2 | 12.9 | 12.2 | 37 |
| 38 | 14.0 | 11.0 | 15.5 | 19.0 | 14.6 | 15.3 | 14.3 | 38 |
| 39 | 14.0 | 12.0 | 17.5 | 20.0 | 16.8 | 17.3 | 15.1 | 39 |
| 40 | 15.0 | 12.5 | 20.0 | 21,0 | 17.6 | 18.8 | 16.3 | 40 |
| 41 | 16.0 | 13.5 | 20.5 | 23.5 | 18.6 | 20.4 | 16.7 | 41 |
| 42 | 17:0 | 14.5 | 24:0 | 25.0 | 19.8 | 20.8 | 18.4 | 42 |
| 43 | 18.0 | 17.5 | 24.5 | 27.0 | 21.2 | 22.0 | 19.2 | +∠ 43 |
| 44 | 20.0 | 20.0 | 26.0 | 29.0 | 22.4 | 22.4 | 20.8 | Same and the same of the same |
| 45 | 21.5 | 23.5 | 28.0 | 32.5 | 23.6 | 23.5 | 20.8 | 44 46 |
| 16 | 24.5 | 23.5 | 28.0 | 33.5 | 26.4 | 26.3 | 24.9 | 45 46 |

Table C.9. (Continued)

| T | <u> </u> | Age-Base | d Sample | | Age-S | Stratified S | ample | т |
|----------|----------|----------|----------|------|---------|--------------|-------|-------|
| Score | 18–24 | 25–44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| 47 | 29.0 | 25.5 | 29.0 | 35.5 | 29.0 | 28.6 | 28.6 | 47 |
| 48 | 31.0 | 28.0 | 31.0 | 36.5 | 31.8 | 32.2 | 30.2 | 48 |
| 49 | 35.5 | 32.5 | 34.0 | 36.5 | 36.2 | 36.5 | 33.1 | 49 |
| 50 | 35.5 | 36.5 | 36.5 | 39.0 | 39.8 | 40.0 | 37.1 | 50 |
| 51 | 39.0 | 41.5 | 42.0 | 45.0 | 42.6 | 44.3 | 40.4 | 51 |
| 52 | 44.0 | 44.0 | 45.5 | 47.5 | 49.2 | 44.3 | 43.7 | 52 |
| 53 | 53.0 | 51.0 | 47.0 | 50.5 | 54.2 | 49.4 | 51.0 | 53 |
| 54 | 58.0 | 56.5 | 52.5 | 55.5 | 59.8 | 54.5 | 57.1 | 54 |
| 55 | 64.0 | 66.5 | 63.5 | 63.0 | 67.0 | 58.4 | 63.7 | 55 |
| 56 | 72.5 | 70.0 | 69.5 | 69.5 | 72.2 | 66.7 | 70.2 | 56 |
| 57 | 79.5 | 79.0 | 78.0 | 72.5 | 78.0 | 72.2 | 74.7 | 57 |
| 58 | 89.5 | 88.5 | 82.0 | 77.0 | 85.6 | 81.2 | 79.2 | 58 |
| 59 | 99.9 | 93.5 | 88.5 | 81.5 | 91.4 | 88.2 | 89.4 | 59 |
| 60 | 99.9 | 99.9 | 95.0 | 84.5 | 95.2 | 92.9 | 95.5 | 60 |
| 61 | 99.9 | 99.9 | 99.9 | 87.0 | 99.9 | 94.9 | 99.9 | 61 |
| -62 | 99.9 | 99.9 | 99.9 | 91.5 | 99.9 | 99,9 | 99.9 | 62 |
| 63 | 99.9 | 99.9 | 99.9 | 95.5 | 99.9 | 99.9 | 99.9 | 63 |
| 64 | 99.9 | 99.9 | 99.9 | 98.0 | 99.9 | 99.9 | 99.9 | 64 |
| ≥65 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | ≥65 |

Table C.10. T Scores Obtained by Cumulative Percentages of the Normative Samples: Mental Health Composite (MHC)

| Τ | | Age-Base | d Sample | | Age-S | Stratified S | ample | Т |
|--|-------------|----------|----------|------|---------|--------------|-------|-------|
| Score | 18-24 | 25–44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| ≤12 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≤12 |
| 13 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≟0:4 | 13 |
| !4 | <0.1 | <0.1 | <0.1 | <0.1 | 0.2 | <0.1 | 0.4 | 14 |
| 15 | <0.1 | <0.1 | <0.1 | <0.1 | 0.4 | <0.1 | 0:4 | 15 |
| 16 | <0.1 | <0.1 | <0.1 | <0.1 | 0.6 | 8.0 | 0.4 | 16 |
| 17 | 0.5 | ≺0.1 | <0.I | <0.1 | 0.6 | 8.0 | .0.4 | 17 |
| 18 | 0.5 | 0.5 | 0.5 | <0.1 | 0.8 | 0.8 | 0.8 | 18 |
| . 19 | 0.5 | 1.0 | 0.5 | <0.1 | 1.2 | 1.2 | 1.2 | 19 |
| 20 | 1.0 | 1.0 | 0.5 | <0.1 | 1.2 | 1.6 | 1.2 | 20 |
| . 21 | 1.5 | 1.5 | 1,0 | <0.1 | 1:4 | 1,6 | 1.2 | 21 |
| 22 | 1.5 | 2.5 | 1.0 | 0.5 | 1.4 | 1.6 | 2.0 | 22 |
| 23 | 2.5 | 2.5 | 1.5 | 1.5 | 2.0 | 2.0 | 2.0 | 23 |
| 24 | 3.0 | 3.0 | 1.5 | 1.5 | 2.6 | 2.7 | 2.9 | 24 |
| 25 | 3.0 | 3.0 | 2.0 | 2.5 | 2.8 | 3.1 | 3:3 | 25 |
| 26 | 4.0 | 3.0 | 3.0 | 3.0 | 3.0 | 3.1 | 3.3 | 26 |
| 27 | 4.0 | 3.5 | 5.0 | 3.0 | 3;4 | 3.1 | 4.1 | 27 |
| 28 | 4.5 | 3.5 | 5.0 | 4.0 | 3.6 | 3.5 | 4.5 | 28 |
| 29 | 6.0 | 3.5 | 5.0 | 4.0 | -4.8 | 4.3 | 4.9 | 29 |
| 30 | 6.0 | 4.0 | 5.5 | 4.0 | 5.4 | 5.5 | 5.3 | 30 |
| 31 | *6.5 - • | 6.0 | 6.5 | 6.5 | 6.2 | 6.3 | 6.1 | 31 |
| 32 | 7.0 | 6.0 | 8.0 | 7.5 | 6.6 | 7.5 | 6.1 | 32 |
| .33 | 8.0 | 7:5 | 8.5 | 8.5 | 7.0 | 7.5 | 7.8 | 33 |
| 34 35 | 8.5 | 8.5 | 9.5 | 10.0 | 8.0 | 7.8 | 7.8 | 34 |
| 36 | 9.5 | 9.0 | 10.5 | 12.0 | 10.4 | 9.4 | 9.4 | 35 |
| STANIS CONTRACTOR CONT | 11.0 | 10.5 | 12.5 | 12.5 | 12.0 | 12.2 | 11.8 | 36 |
| 37 | 12.5 | 11.5 | 13.5 | 13.5 | 13.6 | 12.9 | 13.9 | 37 |
| 38 | 14.5 | 14.5 | 16.0 | 14.5 | 14.4 | 14.5 | 14.3 | 38 |
| 39 | 15.0 | 17.0 | 17.0 | 18.0 | 15.8 | 15,7 | 15.9 | 39 |
| 40 41 | 16.5 | 17.5 | 17.5 | 19.0 | 17.0 | 17.3 | 17.6 | 40 |
| | 17.0 | 19.5 | 20.0 | 19.0 | 18.6 | 18,4 | 18.8 | ·41 |
| 42 | 17.0 | 20.5 | 21.5 | 21.0 | 19.8 | 18.8 | 20.0 | 42 |
| 43 44 | 19.5 | 23.5 | 22.5 | 23.0 | 21.2 | 20.8 | 21.6 | 43 |
| 45 | 22.5 | 24.0 | 24.5 | 24.5 | 23.8 | 22.4 | 23.3 | 44 |
| | 27.0 | 27.0 | 26.0 | 26.0 | 25.8 | 25.9 | 24.1 | 45 |
| 46 47 | 30.0 | 30.0 | 28.0 | 29.0 | 28.6 | 29.8 | 26.5 | 46 |
| 47 | 33.5 | 34:0 | 31.0 | 32.5 | 32.4 | 34.5 | 31.0 | 47 |
| 48 | 35.5 | 37.0 | 36.5 | 36.0 | 36.6 | 37.6 | 34.3 | 48 |
| 49 | 42.0 | 40.5 | 38.5 | 38.0 | 39.4 | 41.2 | 37.1 | 49 |
| 50 | 45.0 | 45.0 | 43.0 | 41.0 | 41.6 | 43.5 | 39.6 | 50 |

Table C.10. (Continued)

| T | | Age-Based | d Sample | | Age-S | Stratified S | ample | T | |
|-------|-------|-----------|-------------------|------|---------|--------------|------------------|----------|--|
| Score | 18–24 | 25–44 | 4564 | ≥65 | Overall | Female | Male | Score | |
| .51 | 46.5 | 46.5 | 46.5 | 47.0 | 44.0 | -46.7 | 42.9 | 51 | |
| 52 | 51.5 | 48.5 | 50.5 | 49.5 | 48.6 | 48.6 | 48.2 | 52 | |
| 53 | 57.5 | 53.0 | 5 4 .5 | 56.5 | 53.2 | .53.7 | .55.1 | .53 | |
| 54 | 61.0 | 58.5 | 60.0 | 64.5 | 58.8 | 57.3 | 62.4 | 54 | |
| 55 | 66.0 | 65.5 | 163.5 | 70.5 | 65.6 | 64.7 | 67.3 | 55 | |
| 56 | 70.5 | 72.0 | 65.5 | 74.5 | 71.2 | 69.8 | 71.4 | 56 | |
| 57 | 76.5 | 76.0 | 74.5 | 77.5 | 75.0 | 72.9 | 75.9 | 57 | |
| 58 | 79.0 | 81.0 | 82.0 | 81.5 | 79.0 | 78.0 | 83.3 | 58 | |
| 59 | 84.5 | 84.5 | 89.0 | 86.5 | 87.2 | 84.3 | 89.4 | 59 | |
| 60 | 90.5 | 88.5 | 90.5 | 86.5 | 92.2 | 89.4 | 91.8 | 60 | |
| 61 | 93.5 | 94.5 | 93.0 | 91.0 | 93.8 | 93.7 | 9 5.1 | -61 | |
| 62 | 95.0 | 95.5 | 96.0 | 93.5 | 96.2 | 94.9 | 95.9 | 62 | |
| 63 | 95.5 | 98.0 | 97.0 | 94.0 | 97.2 | 96.9 | 98.0 | 63 | |
| 64 | 96.5 | 99.0 | 98.0 | 97.0 | 99.0 | 98.4 | 99.2 | 64 | |
| 65 | 98.0 | *99.5 | 99.9 | 98.0 | 99.2 | 99.2 | 99.9 | 65 | |
| 66 | 98.5 | 99.5 | 99.9 | 99.9 | 99.9 | 99.2 | 99.9 | 66 | |
| 67 | 98.5 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 9 9.9 | 67 | |
| ≥68 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | ≥68 | |

Table C.II. T Scores Obtained by Cumulative Percentages of the Normative Samples: Global Health Composite (GHC)

| T | | Age-Base | d Sample | | Age-S | Stratified S | ample | т |
|-------|-------|----------|-----------------|------------------|---------|--------------|--------------|--------|
| Score | 18–24 | 25-44 | 45-64 | ≥65 | 0verall | Female | Male | Score |
| ≤10 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | ≤10 |
| 1.1 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | -11 |
| 12 | 0.5 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 12 |
| 13 | 0.5 | <0.1 | < 0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 13 |
| 14 | 1.5 | 1.0 | <0.1 | <0.1 | 0.4 | <0.1 | <0.1 | 14 |
| 15 | 1.5 | 1:0 | <0.1 | <0.1 | 0.6 | 0.8 | <0.1 | 15 |
| 16 | 1.5 | 1.0 | <0.1 | <0.1 | 0.8 | 1.2 | 0.4 | 16 |
| 17 | 1.5 | 2.0 | <0.1 | <0.1 | 1.0 | 1.2 | 0.8 | 17 |
| 18 | 1.5 | 2.0 | <0.1 | <0.1 | 1.0 | 1.2 | 0.8 | 18 |
| 19 | 1:5 | 2,0 | -<0.1 | ≺ 0.1 | 1.2 | 1.2 | 0,8 | 19 |
| 20 | 1.5 | 2.5 | 1.0 | <0.1 | 1.6 | 1.6 | 1.6 | 20 |
| 21 | 1.5 | 2.5 | 1.0 | <0.1 | 2.0 | 1:6 | 2.0 | 21 |
| 22 | 2.0 | 3.0 | 1.5 | <0.1 | 2.0 | 2.0 | 2.0 | 22 |
| 23 | 2.0 | 3.0 | 1.5 | <0,1 | 2.2 | 2.0 | 2:4 | 23 |
| 24 | 2.0 | 3.0 | 2.0 | <0.1 | 2.8 | 2.0 | 2.9 | 24 |
| 25 | 2.5 | 3.0 | 2.0 | 0.5 | 3.2 | 2.7 | 3.3 | 25 |
| 26 | 2.5 | 3.5 | 2.5 | 1.0 | 3.4 | 2.7 | 4.1 | 26 |
| 27 | 3.0 | 3.5 | 3.0 | 2.5 | 4.2 | 3.1 | 4.5 | 27 |
| 28 | 3.0 | 4.5 | 3.5 | 3.5 | 4.8 | 4.3 | 4.5 | 28 |
| 29 | 3.0 | 4.5 | 4.5 | 4.5 | 5.2 | 4.3 | 5.7 | 29 |
| 30 | 4.0 | 5.0 | 5.5 | 5.5 | 5.8 | 5.1 | 6.1 | 30 |
| 31 | 6.0 | 6.5 | 6.5 | 7.5 | 6.6 | 5.9 | 6.5 | 31 |
| 32 | 6.0 | 7.5 | 8.0 | 8.5 | 7.8 | 7.8 | 7.8 | 32 |
| 33 | 7.0 | 7.5 | 9.0 | 10.5 | 8.2 | 8.6 | 8.2 | 33 |
| 34 | 9.5 | 8.0 | 10.5 | 10.5 | 8.8 | 9.0 | 9.0 | 34 |
| 35 | 11.5 | 8.5 | 11.5 | 12.0 | 9.4 | 9.4 | 9.4 | 35 |
| 36 | 12.5 | 9.5 | 11.5 | 14.0 | 10.8 | 11.0 | 11.0 | 36 |
| 37 | 13.5 | 10:5 | 13.0 | 15.5 | 12.4 | 12:5 | 11:8 | 37 |
| 38 | 15.0 | 13.0 | I <i>4</i> .5 | 16.5 | 14.0 | 13.7 | 14.3 | 38 |
| .39 | 16.0 | 13.5 | 16.0 | 17.5 | 16.0 | 15.3 | 16.3 | 39 |
| 40 | 16.0 | 16.0 | 17.0 | 19.5 | 17.6 | 17.6 | 17.6 | 40 |
| 41 | 17.5 | 16,5 | 20.5 | 23.0 | 18.6 | 18:4 | 18:4 | 41 |
| 42 | 18.5 | 18.5 | 22.0 | 24.5 | 21.0 | 20.0 | 21.6 | 42 |
| 43 | 21.0 | 21.0 | 23.5 | 26.5 | 22.8 | 22.7 | 22.9 | 43 |
| 44 | 24.0 | 22.5 | 27.5 | 28.0 | 24.2 | 24.3 | 24.5 | 44 |
| 45 | 25.5 | 24.0 | 29.0 | 29 .0 | 26.0 | 26.3 | 24.J 26.I | 45 |
| 46 | 27.0 | 27.0 | 32.0 | 30.0 | 28.8 | 29.4 | 28.6 | 46 |
| -47 | 30.5 | 33.0 | 33.0 | 32.0 | 32.2 | 34.5 | 31.0 | 47 |
| 48 | 34.0 | 36.5 | 35.5 | 37.5 | 34.4 | 36.1 | 32.7 | 48 |

Table C.II. (Continued)

| T | | Age-Based | l Sample | | Age-S | Stratified Sa | ample | |
|----------|-------|-----------|----------|------|---------|---------------|-------|-------|
| Score | 18–24 | 25-44 | 45–64 | ≥65 | Overall | Female | Male | Score |
| 49 | 37.5 | 40.5 | 38.5 | 41.0 | 37.0 | 40.4 | 36.7 | 49 |
| 50 | 41.0 | 44.0 | 40.5 | 43.0 | 41.0 | 43.1 | 40.0 | 50 |
| . 51 | 47.0 | 49.5 | 45.0 | 47.0 | -46.2 | 47.8 | 46.5 | 51 |
| 52 | 52.0 | 55.0 | 49.5 | 52.5 | 50.8 | 52.5 | 52.2 | 52 |
| 53 | 56.5 | 59.5 | 54.0 | 57.0 | 56.6 | 56.5 | 55.1 | 53 |
| 54 | 60.5 | 61.5 | 58.5 | 62.0 | 59.6 | 61.2 | 60.4 | 54 |
| 55 | 70.0 | 65.5 | 63:0 | 68.5 | 65.2 | 65.5 | 66.9 | 55 |
| 56 | 75.5 | 72.0 | 71.0 | 72.0 | 71.2 | 72.2 | 71.8 | 56 |
| 57 | 79.5 | 78.0 | 73.0 | 76.0 | 75:8 | 7.6.5 | 78.0 | 57 |
| 58 | 85.0 | 82.5 | 79.0 | 79.0 | 80.8 | 80.8 | 85.7 | 58 |
| 59 | 85.0 | 87.0 | 86,5 | 82.5 | 87.4 | 85.9 | 91.0 | 59 |
| 60 | 90.0 | 92.0 | 91.0 | 85.0 | 92.2 | 91.4 | 93.5 | 60 |
| .6I | 94.0 | 96:0 | 94.5 | 90.5 | 95.6 | 94.9 | 95.9 | 61 |
| 62 | 95.5 | 98.0 | 95.5 | 94.5 | 97.2 | 98.0 | 97.6 | 62 |
| 63 | 97.0 | 99.0 | 98.0 | 97.0 | 98.6 | 98.0 | 99.6 | 63 |
| 64 | 98.0 | 99.9 | 99.0 | 98.5 | 99.8 | 99.6 | 99.6 | 64 |
| .65 | 98.5 | 99.9 | 99.9 | 99.5 | 99.9 | 99.9 | 99.9 | 65 |
| ≥66 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | 99.9 | ≥66 |

Discrepancy Score Tables

Table D.1. Cumulative Percentages of the Normative Samples Obtaining PHC>MHC T-Score Discrepancies

| Amount of | | Age-Base | d Sample | | Age-S | tratified Sa | ımnle |
|----------------|-------|----------|----------|------|---------|--------------|-------|
| Discrepancy | 18–24 | 25-44 | 45-64 | ≥65 | Overall | Female | Male |
| ≥28 | 0.5 | 0.0 | 1.0 | 0.5 | 0.4 | 0.8 | 0.0 |
| 27 | 0.5 | 0.0 | 1.0 | 0.5 | 0.6 | 1.2 | 0.0 |
| 26 | 0.5 | 0.5 | 1.0 | 0.5 | 1.2 | 2.4 | 0.0 |
| 25 | 0.5 | 0.5 | 1.0 | 0.5 | 1.4 | 2.7 | 0.8 |
| 24 | 0.1 | 1.5 | 1.0 | 0.5 | 1.4 | 2.7 | 0.8 |
| 23 | 1.0 | 1.5 | 1.5 | 0.5 | 1.8 | 2.7 | 0.8 |
| 22 | 1.0 | 3.0 | 2.0 | 1.0 | 2.4 | 3.9 | 0.8 |
| 21 | 1:0 | 3.0 | 2.0 | 1.0 | 2.4 | 4.3 | 0.8 |
| 20 | 1.0 | 3.0 | 2.0 | 1.5 | 2.6 | 4.3 | 1.2 |
| 19 | 1.0 | 4.0 | 3.0 | 1.5 | 3.0 | 4.7 | 1:6 |
| 18 | 1.0 | 4.5 | 3.5 | 1.5 | 3.6 | 5.5 | 2.4 |
| 17 | 2.5 | 4.5 | 4.5 | 1.5 | 4.0 | 5.5 | 3.7 |
| 16 | 4.5 | 5.0 | 4.5 | 3.0 | 4.6 | 5.5 | 4.! |
| 15 | 6.0 | 5.5 | 5.0 | 4.5 | 5.0 | 5.5 | 4.5 |
| 14 | 6.5 | 6.0 | 5.5 | 6.0 | 5.2 | 6.7 | 4.9 |
| 13 | 7.0 | 6.5 | 6.5 | 7.0 | 6.8 | 8.2 | 5.7 |
| 12 | 8.5 | 7.0 | 8.0 | 8.0 | 8.2 | 10.2 | 6.9 |
| 11 | 8.5 | 9.5 | 8.5 | 8.0 | 10.2 | 13.7 | 8:6 |
| 10 | 12.0 | 12.0 | 10.0 | 8.5 | 12.0 | 14.9 | 11.0 |
| 9 | 13.5 | 13.0 | 11.5 | 10.5 | 14.4 | 17.6 | 13.5 |
| 8 | 17.5 | 15.5 | 14.5 | 12.5 | 16.6 | 19.2 | 16.7 |
| 7 | 20.0 | 17.5 | 16.5 | 15.5 | 19.2 | 22,7 | 20.8 |
| 6 5 | 24.5 | 21.0 | 20.0 | 20.0 | 23.0 | 25.1 | 24.9 |
| | 26.0 | 24.0 | 22.0 | 24.5 | 26.4 | 29.0 | 26.9 |
| 4 | 31.0 | 29.0 | 27.0 | 27.5 | 29.4 | 32.5 | 30.6 |
| 3 | 37.5 | 35.5 | 31.5 | 35.0 | 32.4 | 36.1 | 35.9 |
| 2 | 41.5 | 42.5 | 35.5 | 44.0 | 36.8 | 38.8 | 41.2 |
| 1 | 44.0 | 47.5 | 42.5 | 47.0 | 42.2 | 44.3 | 46.9 |
| O _a | 7.0 | 7.5 | 6.0 | 6.0 | 5.8 | 5.5 | 6.2 |
| M | _ | | | | | | |
| Mean | -0.2 | 0.2 | 0.0 | 0.1 | -0.4 | 0.0 | 0.2 |
| SD. | 8.9 | 8.5 | 8.1 | 7.6 | 8.9 | 9.9 | 8.1 |
| Median | 0.0 | 0.0 | -1.0 | 0.0 | -1.0 | -1.0 | 0.0 |

Note. PHC-MHC discrepancies >10 were obtained by 15% or less of individuals within each normative group. PHC-MHC discrepancies >6 are statistically significant at p < .10. This table does not include negative PHC-MHC discrepancies. Mean difference scores, standard deviations, and medians are based on the entire distribution.

^aThis row provides the percentage of individuals in each normative group for whom PHC–MHC composite score discrepancies were equal to zero or <1.

Table D.2. Cumulative Percentages of the Normative Samples Obtaining MHC>PHC T-Score Discrepancies

| Amount of | | Age-Base | d Sample | | Age-S | tratified Sa | mple |
|-------------|----------------|----------|----------|---------------|-------------|--------------|------|
| Discrepancy | 18-24 | 25-44 | 4564 | ≥65 | Overall | Female | Male |
| 26 | 0.5 | 0.0 | 0.0 | 0.0 | 0.2 | 0.0 | 0.4 |
| 25 | 1.0 | 0.0 | 0.0 | 0.0 | 0.6 | 0.4 | 0.4 |
| 24 | 1.5 | 0.0 | 0.0 | 0.0 | 0.6 | 0.8 | 0.8 |
| 23 | (2:0 | 0.0 | 0.5 | 0.0 | 0:8 | .0.8 | 0.8 |
| 22 | 2.0 | 0.0 | 0.5 | 0.0 | 1.2 | 0.8 | 0.8 |
| 21 | 2.0 | 1.0 | 0.5 | 1.0 | 1.2 | 1.2 | 1.6 |
| 20 | 2.0 | 1.0 | 1.0 | 1.0 | 1.6 | 2.0 | 1.6 |
| .19 | 2.5 | i5 | 1.5 | 1.0 | 1.8 | 2.0 | ી.6 |
| 18 | 3.0 | 2.5 | 2.0 | 1.5 | 2.2 | 3.1 | 2.0 |
| 17 | 3.5 | 3:0 | 2.0 | 2.0 | 3.0 | 4.3 | 2.4 |
| 16 | 6.5 | 3.5 | 3.5 | 2.0 | 3.6 | 5.1 | 2.4 |
| 15 | 6.5 | 4.5 | 4:0 | 2.0 | ⊬4:4 | 6.3 | 2.9 |
| 14 | 7.5 | 5.0 | 4.5 | 2.5 | 5.2 | 7.8 | 4.9 |
| 139 | 8.5 | 8.0 | 5.0 | 3.0 | 7.6 | 8.6 | 6.1 |
| 12 | 8.5 | 8.5 | 6.0 | 5.0 | 8.6 | 10.6 | 6.9 |
| 41 | 9:0 | 0.01 | 7.0 | 5.5 | 11.6 | 13.7 | 8.6 |
| 10 | 9.5 | 12.0 | 8.5 | 8.0 | 12.4 | 14.5 | 10.6 |
| 9.5 | #I 3. 5 | 13.5 | 11.0 | 10.0 | 15:4 | 17.6 | 11:4 |
| 8 | 14.5 | 15.5 | 13.5 | 13.5 | 18.4 | 20.8 | 14.3 |
| | 17.5 | 17.5 | 15.0 | 19.5 | 22.0 | 25.5 | 17.6 |
| 6 | 23.0 | 21.0 | 19.5 | 22.5 . | 26.6 | 27.8 | 20.4 |
| 577 | 24.5 | 25.5 | 23.5 | 26.0 | 31.2 | 30:2 | 24.5 |
| 4 | 32.5 | 31.5 | 31.5 | 31.5 | 35.8 | 35.3 | 30.6 |
| | 38.5 | 36.5 | 35.5 | 36.0 | 42.8 | 41.6 | 35.1 |
| 2 | 44.5 | 40.0 | 46.5 | 40.0 | 47.0 | 45.9 | 43.3 |
| | 49,0 | -45.0 | 51.5 | 47.0 | 52.0 | 50.2 | 46.9 |
| On | 7.0 | 7.5 | 6.0 | 6.0 | 5.8 | 5.5 | 6.2 |
| | | | | | | | |
| Mean | 0.2 | -0.2 | 0.0 | 0. I | 0.4 | 0.0 | -0.2 |
| SD | 8.9 | 8.5 | 18.1 | 7.6 | 8.9 | 9.9 | 8.1 |
| Median | 0.0 | 0.0 | 1.0 | 0.0 | 1.0 | 1.0 | 0.0 |

Note: MHC-PHC discrepancies >10 were obtained by 15% or less of individuals within each normative group. MHC-PHC discrepancies >6 are statistically significant at p < .10. This table does not include negative MHC-PHC discrepancies. Mean difference scores, standard deviations, and medians are based on the entire distribution.

^{*} his row provides the percentage of individuals in each normative group for whom MHC-PHC composite score discrepancies were equal to zero or <1.